

REVIEW ARTICLE

Obstetrics

Prevalence of postpartum depression in women amid the COVID-19 pandemic: A systematic review and meta-analysis

Meysam Safi-Keykaleh¹ | Fatemeh Aliakbari² | Hamid Safarpour^{3,4} | Mehdi Safari⁵ | Azadeh Tahernejad⁵ | Hojjat Sheikhbardsiri⁶ | Ali Sahebi³

¹Nahavand School of Allied Medical Sciences, Hamadan University of Medical Sciences, Hamadan, Iran

²Community-Oriented Nursing Midwifery Research, Shahrekord University of Medical Sciences, Shahrekord, Iran

³Non-Communicable Diseases Research Center, Ilam University of Medical Sciences, Ilam, Iran

⁴Department of Nursing, School of Nursing and Midwifery, Ilam University of Medical Sciences, Ilam, Iran

⁵Department of Health in Disasters and Emergencies, School of Public Health and Safety, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁶Health in Disasters and Emergencies Research Center, Institute for Future Studies in Health, Kerman University of Medical Sciences, Kerman, Iran

Correspondence

Ali Sahebi, Non-Communicable Diseases Research Center, Ilam University of Medical Sciences, Ilam, Iran.
Email: ali.sahebi.phd@gmail.com

Abstract

COVID-19 outbreaks appear to be related to exacerbation of psychological problems such as depression and anxiety in high-risk population such as pregnant women and the postpartum period due to stress and life-threatening illnesses. The aim of this study was to evaluate the prevalence of postpartum depression (PPD) during COVID-19. This study protocol is registered in PROSPERO with CRD42021278425 code. Data sources including Google Scholar, ISC, Magiran, Scopus, PubMed, Embase, and Web of Science and reference list of included articles were used to identify related studies. Observational studies that reported the prevalence of PPD in both Persian and English during COVID-19 between January 20, 2020 and August 31, 2021 were included. Data were collected and analyzed with a random effects model for meta-analysis. In this study, 671 initial articles were identified and after removing duplicates, 454 studies were screened and finally 24 studies entered the meta-analysis stage. According to this study results PPD based on Edinburgh Postnatal Depression Scale (EPDS) ≥ 9 , EPDS ≥ 10 , EPDS ≥ 11 , EPDS ≥ 12 , EPDS ≥ 13 , Postpartum Depression Screening Scale-Short Form (PDSS-SF) ≥ 17 and total prevalence was reported 12% (95% confidence interval [CI] = 0.07–17, $I^2 = 97\%$), 27% (95% CI = 15–39, $I^2 = 99\%$), 44% (95% CI = 40–49, $I^2 = 0.0\%$), 27% (95% CI = 0.06–49, $I^2 = 97.4\%$), 28% (95% CI = 18–39, $I^2 = 98.5\%$), 37% (95% CI = 32–42), 28% (95% CI = 23–33, $I^2 = 98.5\%$). The findings of this study showed that the prevalence of PPD in women is relatively high during COVID-19. Therefore, considering the psychological consequences of the COVID-19 pandemic and the importance of pregnancy and the postpartum period in terms of the emergence of mental disorders, especially depression, it seems necessary to perform interventions and physical and psychological support.

KEYWORDS

COVID-19, mental health, postpartum depression, women

1 | INTRODUCTION

COVID-19 outbreaks appear to be exacerbated in all populations due to stress and life-threatening risks.¹ The COVID-19 pandemic

is also a trigger for various types of mental health problems such as depression, anxiety and panic disorder.² Measures such as social distancing, reduction of religious activities, tourism, and schooling during COVID-19 have changed people's lifestyles and

mental health outcomes, disproportionately affecting high-risk populations.^{3,4}

Pregnancy and postpartum are challenging and are associated with disorders such as insomnia, relationship tension, and feelings of isolation, which have caused mood disorders in many women. On the other hand, environmental stressors such as natural disasters can exacerbate mood disorders during pregnancy.⁵ Pregnant women experience high stress during epidemics. On the other hand, increased symptoms of depression and anxiety in pregnant women are associated with an increased risk of preterm delivery, postpartum depression (PPD) and behavioral problems in children. Therefore, pregnant women need social support during pregnancy because these measures reduce PPD.^{6,7} According to studies, COVID-19 has been associated with an increased risk of mental health problems in pregnant and postpartum women.⁸

Postpartum depression disorder is the most common postpartum illness among women.⁹ According to the DSM-5 definition, PPD occurs 4 weeks after delivery while others define it as commencing at any time in the first year after delivery.¹⁰ Postpartum depression threatens not only the health of mothers but also the health of infants, and infants of depressed mothers may have developmental disorders, reduced cognitive function, as well as problems pertaining to social communication with their parents and peers.¹¹

Approximately 10%–16% of women in the postpartum period experience a period of major depression and many of them also suffer from anxiety disorders.¹² The results of a study in China showed that the prevalence of depression in pregnant women and postpartum women was 27.43%, and women who were worried about themselves or their baby developing COVID-19 were more prone to depression.⁸ The results of other studies have shown that women who gave birth during COVID-19 had a higher rate of PPD than other women who gave birth during the non-COVID-19 period.¹³

Studies have shown that several studies have been conducted to investigate the association between COVID-19 and PPD in mothers, but a comprehensive study was not found to examine the prevalence of PPD during COVID-19. Access to comprehensive and complete information about the rate of PPD during COVID-19

can be used as a reference for health system decisions and policies. Therefore, the research team decided to conduct a systematic review and meta-analysis study to assess the prevalence of PPD during COVID-19.

2 | MATERIALS AND METHODS

This systematic review and meta-analysis was done according to PRISMA guideline.¹⁴ This study protocol was registered in PROSPERO with CRD42021278425 code.

2.1 | Search strategy

Information sources including Google Scholar, ISC, Magiran, Scopus, PubMed, Embase, and Web of Science and reference list of imported articles were used to identify related studies. Persian equivalent and English key words include: “Postnatal Depression”, “Post-Partum Depression”, “Post-Partum Depression”, “Postpartum Depression”, “Post-Natal Depression”, “Post Natal Depression”, “2019 novel coronavirus disease”, COVID19, “COVID-19 pandemic”, “SARS-CoV-2 infection”, “COVID-19 virus disease”, “2019 novel coronavirus infection”, “2019-nCoV infection”, “Coronavirus disease 2019”, “2019-nCoV disease”, “COVID-19 virus infection” were used. In order to compile the strategy search, keywords, search fields and operators were used. First, Pubmed syntax search was compiled and then based on Pubmed syntax search, syntax search of other databases was compiled. The searches were conducted in both Persian and English from January 20, 2020 to August 31, 2021. Table 1 lists the search strategies in the database types.

2.2 | Eligible criteria

In this study, inclusion criteria include observational studies that reported the prevalence of PPD in both Persian and English

TABLE 1 Search strategy in database types

Database	Search syntax
PubMed	((“Postnatal Depression” OR “Post-Partum Depression” OR “Post-Partum Depression” OR “Postpartum Depression” OR “Post-Natal Depression” OR “Post Natal Depression”) AND (“2019 novel coronavirus disease” OR COVID19 OR “COVID-19 pandemic” OR “SARS-CoV-2 infection” OR “COVID-19 virus disease” OR “2019 novel coronavirus infection” OR “2019-nCoV infection” OR “Coronavirus disease 2019” OR “2019-nCoV disease” OR “COVID-19 virus infection”))
Scopus	((ALL(“Postnatal Depression”) OR ALL(“Post-Partum Depression”) OR ALL(“Post-Partum Depression”) OR ALL(“Postpartum Depression”) OR ALL(“Post-Natal Depression”) OR ALL(“Post Natal Depression”)) AND (ALL(“2019 novel coronavirus disease”) OR ALL(COVID19) OR ALL(“COVID-19 pandemic”) OR ALL(“SARS-CoV-2 infection”) OR ALL(“COVID-19 virus disease”) OR ALL(“2019 novel coronavirus infection”) OR ALL(“2019-nCoV infection”) OR ALL(“Coronavirus disease 2019”) OR ALL(“2019-nCoV disease”) OR ALL(“COVID-19 virus infection”))))
Web of science	((TS = (“Postnatal Depression”) OR TS = (“Post-Partum Depression”) OR TS = (“Post-Partum Depression”) OR TS = (“Postpartum Depression”) OR TS = (“Post-Natal Depression”) OR TS = (“Post Natal Depression”)) AND (TS = (“2019 novel coronavirus disease”) OR TS = (COVID19) OR TS = (“COVID-19 pandemic”) OR TS = (“SARS-CoV-2 infection”) OR TS = (“COVID-19 virus disease”) OR TS = (“2019 novel coronavirus infection”) OR TS = (“2019-nCoV infection”) OR TS = (“Coronavirus disease 2019”) OR TS = (“2019-nCoV disease”) OR TS = (“COVID-19 virus infection”))))

during COVID-19 between January 20, 2020 and August 31, 2021. Exclusion criteria included mean and standard deviation of PPD, PPD reports among men, pregnant women, during epidemics other than COVID-19, as well as systematic review studies, interventional studies, case reports, and letters to the editor.

2.3 | Selection of studies

In this study, 671 studies were entered into the EndNote 7 software after the search results were completed, and after removing duplicates, titles, and abstracts, 454 studies were screened. In the next step, the two researchers independently studied 46 possible related studies in detail and selected 24 final studies.

2.4 | Quality assessment and data extraction

The quality evaluation of the studies was done by two researchers independently and any disagreement between them was resolved through a third person. To evaluate cross-sectional studies from the checklist Appraisal tool for Cross-Sectional Studies (AXIS) Tool¹⁵ was used. This tool has a score of 0–20. The Newcastle-Ottawa Scale (NOS) tool¹⁶ was also used to evaluate other observational studies

(cohort and case-control). This tool has eight items in cohort studies in three sections including Selection (four items), Comparability (one item) and Outcome (three items) and also in case-control studies eight items including selection (Selection (four items), Comparability (one item) and Exposure (three items). Also, to extract the required data, two researchers independently used a pre-prepared checklist including: first author, study site, sample size, tools, and the prevalence of PPD used.

2.5 | Statistical analysis

In this study, the binomial distribution formula was used to calculate the standard error of prevalence in each study. Random effects model was used to estimate pooled measures and 95% confidence interval [CI]. The I^2 index was used to examine the heterogeneity between studies and heterogeneity less than 25%, 25%–50%, 50%–75% and more than 75% indicate no heterogeneity, moderate heterogeneity, high and very high.¹⁷ The effect of heterogeneity on the prevalence of PPD was investigated by analyzing the subgroups based on the type of tools and also the cut-off points of the tools. Publication bias in the study was assessed using the Begg test. All analyzes were performed using STATA software, version 14 (StataCorp., College Station, TX, USA).




FIGURE 1 Flowchart of the selection of studies based on PRISMA

3 | RESULTS

In this review study, 671 initial articles were identified through search and after removing duplicates, 454 studies were screened and finally 24 studies were selected and evaluated for quality and all of them entered the meta-analysis stage (Figure 1). Also in this study, 13 169 women were examined for postpartum depression during COVID-19. In this study, Edinburgh Postnatal Depression Scale (EPDS) and Postpartum Depression Screening Scale-Short Form (PDSS-SF) were used to evaluate PPD.

Among the selected methodology studies, 18 were cross-sectional studies, five were retrospective cohort studies, and two were case control studies. The characteristics of other studies are also listed in Table 2. Based on the results of this study, three studies were performed based on EPDS ≥ 9 and PPD levels 12% (95% CI = 0.07–17, $I^2 = 97\%$, $P = 0.009$) were reported. Eight studies were done based on EPDS ≥ 10 and PPD 27% (95% CI = 15–39, $I^2 = 99\%$, $P < 0.001$) was reported. Three study was done based on EPDS

≥ 11 and PPD 44% (95% CI = 40–49, $I^2 = 0.0\%$, $P = 0.436$) was reported. Three studies were done based on EPDS ≥ 12 and PPD 27% (95% CI = 0.06–49, $I^2 = 97.4\%$, $P < 0.001$). Seven studies were done based on EPDS ≥ 13 and PPD 28% (95% CI = 18–39, $I^2 = 98.5\%$, $P < 0.001$) was reported. One study was done based on PDSS-SF ≥ 17 and PPD 37% (95% CI = 32–42) was reported. Also, total prevalence of PPD in women during COVID-19, 28% (95% CI = 23–33, $I^2 = 98.5\%$, $P < 0.001$) (Figure 2). Index I^2 showed that heterogeneity between studies is very high (Figure 2). Based on the Begg test results ($P = 0.084$), the publication bias is not significant in the present study (Figure 3).

4 | DISCUSSION

Based on the results of this review study, 24 articles were reviewed and meta-analyzed and the overall prevalence of PPD in the COVID-19 pandemic was reported to be 28%. Also, the results

TABLE 2 Characteristics of entered study in systematic review and meta-analysis

First author	Type of study	Location	Tools	Sample size	Prevalence of PPD (%)
An ¹⁸	Cross-sectional	China	EPDS ≥ 10	209	56.9
Ceulemans ¹⁹	Cross-sectional	Ireland, Norway, Switzerland, Netherlands, United Kingdom	EPDS ≥ 13	5134	13
Stojanov ²⁰	Cross-sectional	Serbia	EPDS ≥ 10	108	14.8
Fallon ²¹	Cross-sectional	UK	EPDS ≥ 13	614	43
Guvenc ²²	Cross-sectional	Turkey	EPDS ≥ 13	212	34.0
Hiiragi ²³	Retrospective cohort	Japan	EPDS ≥ 9	279	14
Baran ¹¹	Cross-sectional	Poland	EPDS ≥ 11	130	40
Liang ²⁴	Cross-sectional	China	EPDS ≥ 10	864	30.0
Loret de Mola ²⁵	Retrospective cohort	Brazil	EPDS ≥ 13	1042	29.3
Mariño-Narvaez ²⁶	Cross-sectional	Spanish	EPDS ≥ 10	75	22.4
Oskovi-Kaplan ²⁷	Cross-sectional	Turkey	EPDS ≥ 13	223	14.7
Ostacoli ²⁸	Cross-sectional	Italy	EPDS ≥ 11	163	44.2
Pariante ²⁹	Retrospective cohort	Israel	EPDS ≥ 10	223	16.7
Spinola ³⁰	Cross-sectional	Italy	EPDS ≥ 12	243	44
Suárez-Rico ³¹	Cross-sectional	Mexico	EPDS ≥ 13	293	39.2
Suzuki ³²	Cross-sectional	Japan	EPDS ≥ 9	132	14.4
Myers ³³	Cross-sectional	United Kingdom	EPDS ≥ 11	162	47.5
Vatcheva ³⁴	Cross-sectional	Belgium	EPDS ≥ 13	34	26
Hui ³⁵	Retrospective cohort	Hong Kong	EPDS ≥ 10	802	2.9
Hui ³⁵	Retrospective cohort	Hong Kong	EPDS ≥ 10	925	14.4
Miranda ⁴	Cross-sectional	Argentina	PDSS-SF ≥ 17	305	37
Terada ³⁶	Cross-sectional	Japan	EPDS ≥ 9	461	7.6
Zanardo ¹³	Case – control	Italy	EPDS ≥ 12	91	26
Tarabay ³⁷	Cross-sectional	Saudi Arabia	EPDS ≥ 10	150	60.7
Madera ³⁸	Case – control	Italy	EPDS ≥ 12	295	11.9

Abbreviations: EPDS, Edinburgh Postnatal Depression Scal; PDSS-SF, Postpartum Depression Screening Scale-Short Form; PPD, postpartum depression.

Bold values indicate statistically significant.

PPD prevalence 25%.⁴¹ The results of another study that predicted PPD in earthquake survivors showed that based on EPDS ≥ 13 , the PPD rate is 13%.⁴² According to studies, COVID-19 causes stress due to factors such as unpredictability, uncertainty about disease control, and serious life-threatening risks.¹ COVID-19 also exacerbates mental illness, feelings of hopelessness, anxiety, and social isolation.⁴³ On the other hand, pregnancy and childbirth have been two important events in a woman's life and the birth of a baby causes sudden and drastic changes in a woman's roles and responsibilities. Therefore, the postpartum period indicates the time of onset of PPD.⁴⁴ In reviewing the results of studies and the present study, it can be concluded that the prevalence of PPD during COVID-19 was higher than other disasters as well as non-COVID-19 conditions, and based on what was said, it can be concluded that mothers during COVID-19, in addition to experiencing important events such as pregnancy and childbirth, are exposed to severe psychological consequences, especially stress and anxiety, which can increase the risk of PPD. Therefore, in order to prevent or reduce the rate of PPD, it is recommended that mothers, as a vulnerable group, receive special physical and psychological care and support during COVID-19, and in case of depressive symptoms, special measures and interventions are needed, as well as frequent follow-ups performed by health centers and hospitals because PPD may affect mothers' social behaviors and may lead to persistent depression.

The COVID-19 outbreak has had significant effects on vulnerable groups, especially pregnant women. Additionally, the pregnancy and postpartum periods are vulnerable periods that can increase the level of distress in many women.⁴⁵ According to other studies, factors such as economic and social factors, history of childbirth, lifestyle and history of mental illness, level of education, gestational diseases, family dissatisfaction and limited communication and interaction with others are important predictors of PPD.^{46,47} Studies show that several factors play a role in the occurrence of PPD, for example, COVID-19 as a stressor can accelerate the occurrence of PPD, so mothers should be under close monitoring during pregnancy and after childbirth by families and health care workers. Also, it is recommended that government officials, psychologists and health managers provide stress management training to identify and diagnose women with a history of mental disorders and develop programs and protocols for mental health support for women during and after pregnancy.

5 | CONCLUSION

The findings of this study showed that the prevalence of PPD in women was relatively high during COVID-19. Since PPD can lead to major depression and dangerous behaviors such as suicide and other life-threatening behaviors and have impacts on social behaviors, considering the psychological consequences of the COVID-19 pandemic in this period is important. Psychological interventions are

necessary in pregnancy and after childbirth for prevention of mental disorders, especially depression. Prevention, reduction, diagnosis, treatment and rehabilitation of PPD require screening, frequent follow-ups, training, interventions and physical and psychological support from health care centers and hospitals. It is also recommended that studies should be undertaken to identify the challenges of pregnant and postpartum women in disasters to provide preventive solutions, formulate strategies and appropriate responses and rehabilitation solutions.

AUTHOR CONTRIBUTIONS

MS-K and AS conceived the study. Meysam Safi-Keykaleh, Hamid Safarpour and Ali Sahebi designed the study. Meysam Safi-Keykaleh, Fatemeh Aliakbari, Mehdi Safari, Hojjat Sheikhbardsiri and Azadeh Tahernejad screened the abstracts for inclusion in the study. Ali Sahebi and Hamid Safarpour analyzed the data. Meysam Safi-Keykaleh and Ali Sahebi drafted the manuscript, which was then critically revised by all authors. All authors approved the final manuscript.

CONFLICTS OF INTEREST

The authors have no conflicts of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

REFERENCES

- Adibi A, Golitaleb M, Farrahi-Ashtiani I, et al. The prevalence of generalized anxiety disorder among health care workers during the COVID-19 pandemic: a systematic review and meta-analysis. *Front Psychiatry*. 2021;12(645):658846.
- Qiu J, Shen B, Zhao M, Wang Z, Xie B, Xu Y. A nationwide survey of psychological distress among Chinese people in the COVID-19 epidemic: implications and policy recommendations. *Gen Psychiatry*. 2020;33(2):e100213.
- Davenport MH, Meyer S, Meah VL, Strynadka MC, Khurana R. Moms are not OK: COVID-19 and maternal mental health. *Front Glob Womens Health*. 2020;1:1.
- Miranda AR, Scotta AV. Triggering of postpartum depression and insomnia with cognitive impairment in Argentinian women during the pandemic COVID-19 social isolation in relation to reproductive and health factors. *Midwifery*. 2021;102:103072.
- Farewell CV, Jewell J, Walls J, Leiferman JA. A mixed-methods pilot study of perinatal risk and resilience during COVID-19. *J Prim Care Community Health*. 2020;11:2150132720944074.
- Jago CA, Singh SS, Moretti F. Coronavirus disease 2019 (COVID-19) and pregnancy: combating isolation to improve outcomes. *Obstet Gynecol*. 2020;136(1):33-36.
- Lebel C, MacKinnon A, Bagshawe M, Tomfohr-Madsen L, Giesbrecht G. Elevated depression and anxiety symptoms among pregnant individuals during the COVID-19 pandemic. *J Affect Disord*. 2020;277:5-13.
- Bo H-X, Yang Y, Chen J, et al. Prevalence of depressive symptoms among pregnant and postpartum women in China during the COVID-19 pandemic. *Psychosom Med*. 2021;83(4):345-350.
- Zlotnick C, Miller IW, Pearlstein T, Howard M, Sweeney P. A preventive intervention for pregnant women on public

- assistance at risk for postpartum depression. *Am J Psychiatry*. 2006;163(8):1443-1445.
10. Hamami H, Sheiner E, Wainstock T, et al. The association between delivery during the COVID-19 pandemic and immediate postpartum maternal cognitive function. *J Clin Med*. 2020;9(11):3727.
 11. Baran J, Leszczak J, Baran R, et al. Prenatal and postnatal anxiety and depression in mothers during the COVID-19 pandemic. *J Clin Med*. 2021;10(14):3193.
 12. Misri S, Reebye P, Corral M, Milis L. The use of paroxetine and cognitive-behavioral therapy in postpartum depression and anxiety: a randomized controlled trial. *J Clin Psychiatry* 2004;65(9):0-1236, 1241.
 13. Zanardo V, Manghina V, Gilberti L, Vettore M, Severino L, Straface G. Psychological impact of COVID-19 quarantine measures in northeastern Italy on mothers in the immediate postpartum period. *Int J Gynecol Obstet*. 2020;150(2):184-188.
 14. Moher D, Liberati A, Tetzlaff J, Altman DG, for the PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(7):e1000097.
 15. Downes MJ, Brennan ML, Williams HC, Dean RS. Development of a critical appraisal tool to assess the quality of cross-sectional studies (AXIS). *BMJ Open*. 2016;6(12):e011458.
 16. Wells GA, Shea B, O'Connell D, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. 2014.
 17. Sahebi A, Abdi K, Moayedi S, Torres M, Golitaleb M. The prevalence of insomnia among health care workers amid the COVID-19 pandemic: an umbrella review of meta-analyses. *J Psychosom Res*. 2021;149:110597.
 18. An R, Chen X, Wu Y, et al. A survey of postpartum depression and health care needs among Chinese postpartum women during the pandemic of COVID-19. *Arch Psychiatr Nurs*. 2021;35(2):172-177.
 19. Ceulemans M, Foulon V, Ngo E, et al. Mental health status of pregnant and breastfeeding women during the COVID-19 pandemic—a multinational cross-sectional study. *Acta Obstet Gynecol Scand*. 2021;100:1219-1229.
 20. Stojanov J, Stankovic M, Zikic O, Stankovic M, Stojanov A. The risk for nonpsychotic postpartum mood and anxiety disorders during the COVID-19 pandemic. *Int J Psychiatry Med*. 2021;56(4):228-239.
 21. Fallon V, Davies SM, Silverio SA, Jackson L, De Pascalis L, Harrold JA. Psychosocial experiences of postnatal women during the COVID-19 pandemic. A UK-wide study of prevalence rates and risk factors for clinically relevant depression and anxiety. *J Psychiatr Res*. 2021;136:157-166.
 22. Guvenc G, Yesilcinar İ, Ozkececi F, et al. Anxiety, depression, and knowledge level in postpartum women during the COVID-19 pandemic. *Perspect Psychiatr Care*. 2021;57(3):1449-1458.
 23. Hiiragi K, Obata S, Misumi T, Miyagi E, Aoki S. Psychological stress associated with the COVID-19 pandemic in postpartum women in Yokohama, Japan. *J Obstet Gynaecol Res*. 2021;47(6):2126-2130.
 24. Liang P, Wang Y, Shi S, Liu Y, Xiong R. Prevalence and factors associated with postpartum depression during the COVID-19 pandemic among women in Guangzhou, China: a cross-sectional study. *BMC Psychiatry*. 2020;20(1):1-8.
 25. Loret de Mola C, Martins-Silva T, Carpena MX, et al. Maternal mental health before and during the COVID-19 pandemic in the 2019 Rio Grande birth cohort. *Braz J Psychiatry*. 2021;43:402-406.
 26. Mariño-Narvaez C, Puertas-Gonzalez JA, Romero-Gonzalez B, Peralta-Ramirez MI. Giving birth during the COVID-19 pandemic: the impact on birth satisfaction and postpartum depression. *Int J Gynecol Obstet*. 2021;153(1):83-88.
 27. Oskovi-Kaplan ZA, Buyuk GN, Ozgu-Erdinc AS, Keskin HL, Ozbas A, Tekin OM. The effect of COVID-19 pandemic and social restrictions on depression rates and maternal attachment in immediate postpartum women: a preliminary study. *Psychiatr Q*. 2021;92(2):675-682.
 28. Ostacoli L, Cosma S, Bevilacqua F, et al. Psychosocial factors associated with postpartum psychological distress during the COVID-19 pandemic: a cross-sectional study. *BMC Pregn Childbirth*. 2020;20(1):1-8.
 29. Pariente G, Broder OW, Sheiner E, et al. Risk for probable postpartum depression among women during the COVID-19 pandemic. *Arch Womens Ment Health*. 2020;23(6):767-773.
 30. Spinola O, Liotti M, Speranza AM, Tambelli R. Effects of COVID-19 epidemic lockdown on postpartum depressive symptoms in a sample of Italian mothers. *Front Psychiatry*. 2020;11:1177.
 31. Suárez-Rico BV, Estrada-Gutierrez G, Sánchez-Martínez M, et al. Prevalence of depression, anxiety, and perceived stress in postpartum Mexican women during the COVID-19 lockdown. *Int J Environ Res Public Health*. 2021;18(9):4627.
 32. Suzuki S. Psychological status of postpartum women under the COVID-19 pandemic in Japan. *J Matern Fetal Neonatal Med*. 2020;1-3.
 33. Myers S, Emmott EH. Communication across maternal social networks during England's first national lockdown and its association with postnatal depressive symptoms. *Front Psychol*. 2021;12:1583.
 34. Vatcheva T, Mostaert A, Van Ingelgem V, Henrion E, Legros L. Impact of COVID-19 pandemic on postpartum depression among mothers of extreme and early preterm infants. *Int J Gynecol Obstet*. 2021;155:490-495.
 35. Hui P, Ma G, Seto T, Cheung K. Effect of COVID-19 on delivery plans and postnatal depression scores of pregnant women. *Hong Kong Med J*. 2020;27:113-117.
 36. Terada S, Kinjo K, Fukuda Y. The relationship between postpartum depression and social support during the COVID-19 pandemic: a cross-sectional study. *Journal of Obstetrics and Gynaecology Research*. 2021;47:3524-3531.
 37. Tarabay AI, Boogis D, Tabbakh AT, et al. Prevalence and factors associated with postpartum depression during the COVID-19 Pandemic among Women in Jeddah, Saudi Arabia: a cross-sectional study. *Open Journal of Obstetrics and Gynecology*. 2020;10(11):1644-1657.
 38. Madera P, Romagnolo C, Zanetti S, Dal Maso F, Turro M, De Marco C. Effects of the COVID-19 pandemic on postpartum depression: a retrospective case-control study on a significant sample of mothers in northern Italy. *J Prenat Perinat Psychol Health*. 2021;35:589916.
 39. Yan H, Ding Y, Guo W. Mental health of pregnant and postpartum women during the coronavirus disease 2019 pandemic: a systematic review and meta-analysis. *Front Psychol*. 2020;11:3324.
 40. Demissie DB, Bitew ZW. Mental health effect of COVID-19 pandemic among women who are pregnant and/or lactating: a systematic review and meta-analysis. *SAGE Open Med*. 2021;9:20503121211026195.
 41. Alshikh Ahmad H, Alkhatib A, Luo J. Prevalence and risk factors of postpartum depression in the Middle East: a systematic review and meta-analysis. *BMC Pregn Childbirth*. 2021;21(1):1-12.
 42. Demirchyan A, Petrosyan D, Armenian HK. Rate and predictors of postpartum depression in a 22-year follow-up of a cohort of earthquake survivors in Armenia. *Arch Womens Ment Health*. 2014;17(3):229-237.
 43. Sahebi A, Nejati-Zarnaqi B, Moayedi S, Yousefi K, Torres M, Golitaleb M. The prevalence of anxiety and depression among healthcare workers during the COVID-19 pandemic: an umbrella review of meta-analyses. *Prog Neuropsychopharmacol Biol Psychiatry*. 2021;107:110247.
 44. Slomian J, Honvo G, Emonts P, Reginster J-Y, Bruyère O. Consequences of maternal postpartum depression: a systematic review of maternal and infant outcomes. *Womens Health*. 2019;15:1745506519844044.

45. Boekhorst MG, Muskens L, Hulsbosch LP, et al. The COVID-19 outbreak increases maternal stress during pregnancy, but not the risk for postpartum depression. *Arch Womens Ment Health*. 2021;24(6):1037-1043.
46. Do TKL, Nguyen TTH, Pham TTH. Postpartum depression and risk factors among Vietnamese women. *Biomed Res Int*. 2018;2018:1-5.
47. Ghaedrahmati M, Kazemi A, Kheirabadi G, Ebrahimi A, Bahrami M. Postpartum depression risk factors: a narrative review. *J Educ Health Promot*. 2017;6:60.

How to cite this article: Safi-Keykaleh M, Aliakbari F, Safarpour H, et al. Prevalence of postpartum depression in women amid the COVID-19 pandemic: A systematic review and meta-analysis. *Int J Gynecol Obstet*. 2022;157:240–247. doi: [10.1002/ijgo.14129](https://doi.org/10.1002/ijgo.14129)